

# BRITISH PARACHUTE ASSOCIATION LTD

5 Wharf Way, Glen Parva, Leicester, LE2 9TF

Telephone: 0116 278 5271, Fax: 0116 247 7662, e-mail: [skydive@bpa.org.uk](mailto:skydive@bpa.org.uk)



[www.bpa.org.uk](http://www.bpa.org.uk)

**(This form should not be sent to the British Parachute Association)**

## DECLARATION OF FITNESS TO PARACHUTE

I hereby declare that I am physically fit. I do not, and have not, suffered from any of the following conditions, which I understand may lead to a dangerous situation with regard to myself or other persons during parachuting:

Epilepsy, fits, severe head injury, recurrent blackouts or giddiness, disease of the brain or nervous system, high blood pressure, heart or lung disease, recurrent weakness or dislocation of any limb, diabetes, mental illness, drug or alcohol addiction.

I further declare that in the event of contracting or suspecting any of the above conditions, or in the event of sickness absence over twenty consecutive days, incapacitating injury or confirmation of pregnancy, I will cease to parachute until I have obtained medical approval. I have read the notes overleaf.

.....  
Name in CAPITALS

.....  
Date of Birth

.....  
Weight

.....  
Signature

.....  
Date

.....  
BPA Number

.....  
Height

(BPA Number may be issued on day of Course)

.....  
Signature of Witness

.....  
Name of Witness in CAPITALS

(All parachutists require their Declaration Witnessed, for parachutists under 18 years of age, the Witness MUST be the parent or guardian)

.....  
IF YOU CANNOT SIGN THE DECLARATION BECAUSE OF ANY OF THE ABOVE CONDITIONS, OR IF YOU ARE AGED 40 OR MORE, YOU MUST OBTAIN THE DOCTORS CERTIFICATE BELOW BEFORE PARACHUTING. THIS IS NOT N.H.S. WORK AND YOUR DOCTOR MAY CHARGE YOU FOR THIS.  
.....


## DOCTOR'S CERTIFICATE

I understand that the applicant wishes to parachute but is unable to sign the above declaration/aged 40 or over\*. I have read the notes overleaf. In my opinion the applicant is physically and mentally capable of parachuting and is safe to do so. Glasses or contact lenses must/need not\* be worn.

.....  
Signature

.....  
Date of Signature

.....  
Date of Expiry  
(see – Validity, over)



(Doctors Stamp)

\* Delete as applicable

**NOTES FOR PARACHUTISTS** Parachutists need a reasonably high standard of physical fitness and must not be overweight in relation to their sex, age, and height. No person under the age of 16, or over the age of 55 years will be permitted to carry out initial 'solo' parachute training (see BPA Form 114A – 'Solo' Student Parachutist Declaration of Fitness to Parachute/Doctor's certificate). Exceptions to the higher age limit may be permitted if the person has previous recorded parachute experience. Higher age limits for Student Tandem Parachutists may be acceptable (see BPA Form 115 – Student Tandem Parachutist Declaration of Fitness to Parachute/Doctor's certificate).

As well as the conditions listed overleaf, the following may cause problems to parachutists and if you have ever suffered from any of them you must seek medical approval before parachuting:

Previous fractures, back strain, arthritis and severe joint sprains. Chronic bronchitis. Asthma. Rheumatic fever. Pneumothorax. Liver or Kidney disease. Thyroid, adrenal or other glandular disorder. Chronic ear or sinus disease. Any condition which requires the regular use of drugs. Anaemia. Recent blood donation.

If you wear spectacles they should be securely attached while parachuting. If contact lenses are used, protective goggles should be worn. Your sight must be adequate to read a car number plate at 25 yards.

**NOTES FOR DOCTORS** Cardiorespiratory fitness is important. Sport parachutists make descents from unpressurised aircraft at heights of 2,000 to 15,500 feet above sea level without supplementary oxygen. At 15,000 feet there is a 40% reduction in available oxygen. A tachycardia of 120 - 160 bpm is common in experienced parachutists and 200 bpm is not unusual in novices. The tachycardia may be present at the same time as relative hypoxia and considerable physical exertion.

Musculoskeletal fitness is required. The parachutist must be able to operate his equipment with either hand and should be able to exert a pull of 30 lbs with either hand in any direction. During the parachute deployment there is a brisk deceleration, usually about 4g but occasionally up to 15g. The landing impact typically involves a descent rate equivalent to jumping from a wall 4 feet high, with a horizontal speed of 0 - 15mph. Occasionally the landing impact may be considerably greater than this. Pre-existing spinal or joint injuries may be exacerbated. Obesity increases the likelihood of lower limb or spinal injuries.

A visual acuity of at least 6/12 (after correction with spectacles or contact lenses) is required. Blindness in one eye is acceptable provided that the remaining eye has a full field and the candidate has adapted to monocular vision. Middle ear or sinus disease may cause problems due to the rapid changes in ambient air pressure. The rate of descent in freefall may exceed 10,000 ft/min and under an open canopy 1,000 ft/min. Normal peripheral sensation and coordination are required to activate the parachute but may be impaired by disorders of the nervous system or peripheral vasculature. On early jumps the candidate is responsible mainly for his own safety but must behave in such a way as to cause no hazard to others. After further training, but no further medical evidence, the candidate may be the sole person responsible for the safety of an aircraft full of student parachutists. The candidate must be physically and psychologically fit to carry this responsibility.

The following conditions will normally make a candidate UNFIT to parachute, although there are some exceptions: Any condition which can lead to blackouts, impaired consciousness or impaired concentration. Tendency to persistent or recurrent weakness or dislocation of any limb (unless successfully corrected by surgery). Conditions requiring the use of medication with sedative or psychotropic side effects. A history of ischaemic heart disease, uncontrolled hypertension, other significant cardiac or respiratory disease, cerebrovascular disease, epilepsy, diabetes, mental illness, drug addiction, alcohol dependence, significant CNS disease. A certifying doctor is not stating that a candidate will remain free of injury during parachuting, but that records, history or appropriate clinical examination have not suggested unacceptable medical risk factors. In cases of doubt, or where further information is required, the Medical Adviser to the British Parachute Association or the National Coach and Safety Officer will be pleased to help, and may be contacted at the address overleaf.

These notes are not exhaustive. Some candidates who are 'unfit' using the above criteria may still be fit to make a tandem parachute descent (strapped into a common harness with an experienced instructor) - see separate 'Student Tandem Parachutist' form.

**VALIDITY** The *Parachutist's Declaration* overleaf places the parachutist under a permanent obligation to cease parachuting until obtaining a doctors approval if he/she develops any of the listed conditions. It is valid initially up to the age of 40, but should be renewed every 2 years between the ages of 40 and 49 and every year from the age of 50. *Doctor's Certificates* are valid as follows:

- |                          |   |  |
|--------------------------|---|--|
| If required under age 40 | - | As indicated by stability of medical condition (but not beyond age 40) |
| Age 40 - 49 years        | - | The longer of "3 years" or "until age 50"                              |
| Age 50 years or over     | - | 3 years  |

providing that the examining doctor can specify a shorter period of validity if he/she feels it appropriate.